

**AT** MICROFICHE  
REFERENCE  
LIBRARY

A project of Volunteers in Asia

A Farmer's Primer on Growing Rice

by Benito S. Vergara

Published by:

International Rice Research Institute  
P.O. Box 933  
Manila  
PHILIPPINES

Available from:

same as above

Reproduced by permission.

Reproduction of this microfiche document in any form is subject to the same restrictions as those of the original document.

# **A FARMER'S PRIMER ON GROWING RICE**


**Benito S. Vergara**

1979

**INTERNATIONAL RICE RESEARCH INSTITUTE**

LOS BAÑOS, LAGUNA, PHILIPPINES

P.O. BOX 933, MANILA, PHILIPPINES



# Contents

- 1 Life cycle of the rice plant
- 9 The seed
- 19 Seedling growth
- 29 How to select good seedlings
- 37 Transplanting
- 43 The leaves
- 49 The roots
- 65 The tillers
- 77 The panicle
- 85 Dormancy
- 91 Fertilizers
- 99 How much nitrogen to apply
- 107 How to increase the efficiency of  
nitrogen fertilizer
- 117 Why more nitrogen fertilizer is applied  
during the dry season
- 123 Carbohydrates production
- 133 Water
- 141 Yield components
- 155 Plant type of a lowland rice variety  
with high grain potential
- 167 Factors affecting lodging
- 177 Weeds
- 189 Control of weeds
- 197 Herbicides
- 209 How to judge a rice crop at flowering

# Foreword

A PROGRESSIVE RICE FARMER should understand *why* and *how* the improved rice varieties and farm technology increase production. But recommendations given to farmers often fail to answer questions such as why a farmer incubates seed, why he or she applies fertilizer, or how and when that fertilizer should be incorporated.

The farmer needs this knowledge to adjust his practices to suit his own unique farm situation.

To improve the understanding of rice culture among farmers, technicians, teachers, and scientists, B. S. Vergara of the IRRI plant physiology department prepared this handbook. *A farmer's primer on growing rice* should be particularly useful to technicians and the farmers that they serve. Dr. Vergara initiated work on the handbook during a sabbatic leave at the Southeast Asian Regional Center for Graduate Study and Research in Agriculture (SEARCA), Los Baños, Philippines. Donald Esslinger, editor of the Missouri State Agricultural Extension Service, took the leadership in editing *A farmer's primer on growing rice* while on sabbatic leave with IRRI's Office of Information Services (OIS).

**N. C. Brady**

Director General


International Rice Research Institute

# LIFE CYCLE OF THE RICE PLANT

- 3 The rice plant
- 4 Growth stages of the rice plant
- 5 Difference in growth stages
- 6 Vegetative phase
- 7 Productive phase
- 8 Ripening phase


# THE RICE PLANT

RICE WITH FIVE TILLERS



- A tiller is a shoot which includes the roots, stem, and leaves. It may or may not have a panicle.

# GROWTH STAGES OF THE RICE PLANT





Seeding	Transplanting	Maximum tiller number	Panicle formation	Flowering	Harvest
Vegetative Phase 55 days			Reproductive Phase 35 days	Ripening Phase 30 days	


- Days in vegetative phase differ with variety.
- Reproductive and ripening phases are constant for most varieties. Panicle formation to flowering is 35 days. Flowering to harvest requires 30 days.
- Sowing to harvest may be 180 days or more.




# DIFFERENCE IN GROWTH STAGES



- Number of days in vegetative phase changes by variety.
- Days in reproductive and ripening phases are more or less fixed.
- Difference in growth duration is determined by days in vegetative phase.




Seedling Stage



Tillering Stage

- Duration of the seedling or nursery stages:
  - Dapog (9–11 days)
  - Wetbed (16–20 days)
  - Direct seeding (none)
- Number of tillers and leaf area increase during the vegetative stage.
- Low temperature or long day length can increase the duration of the vegetative phase.

## PRODUCTIVE PHASE



- The reproductive phase begins at the start of panicle formation and ends at flowering. This takes around 35 days.

## RIPENING PHASE



- The ripening phase starts at flowering and lasts for 30 days.
- Rainy days or low temperatures may delay the ripening phase, while sunny and warm days shorten it.
- To obtain high grain yields, good farming practices are needed at each growth stage.

# THE SEED


- 11 The seed
- 12 Parts of a seed
- 13 Stages of germination
- 14 Conditions needed for seed germination — water
- 15 Conditions needed for seed germination — air
- 16 Conditions needed for seed germination — warm temperature
- 17 Why incubate the seeds
- 18 Why select good seeds

## THE SEED



● Seeds vary in size, shape, color, and length of awn.


## PARTS OF A SEED



The seed was cut lengthwise

- The hull is the hard cover of the seed.
- The endosperm is made up mostly of starch, sugar, protein and fats. It is the storehouse of food for the embryo.
- Almost 80 percent of the endosperm is starch. The food needed for seed germination is in the endosperm.
- The embryo develops into the shoot and the roots.


# STAGES OF GERMINATION



- Growth of the embryo will depend on temperature and availability of water and air.




# CONDITIONS NEEDED FOR SEED GERMINATION – WATER




- Uptake of water is the first need of a seed for germination.
- There are many activities going on inside the germinating seed. Starch, protein and fats are being changed into simple foods for the embryo.
- Soak seeds for at least 24 hours so that water can easily and uniformly enter the seeds.

## CONDITIONS NEEDED FOR SEED GERMINATION – AIR




- The germinating rice seed needs air to live.
- Water contains very little air.
- If the seed is covered too deeply with water, the growth of the embryo will be slow and the resulting shoot is tall and weak.

# CONDITIONS NEEDED FOR SEED GERMINATION—WARM TEMPERATURE




- Warm temperature is needed to increase the activities inside the seed and thus increase growth.
- Low temperature decreases activities inside the seed.

# WHY INCUBATE THE SEEDS



Soak 24 hours



Incubate 24 hours

After soaking for 24 hours, water is removed and seeds are washed, placed on a cement floor and covered with a sack.

- Incubation keeps the seeds warm, increases growth of the embryo, and results in uniform germination.
- If the incubation temperature is too high, germination rate decreases and may actually kill the sprouted seeds.

## WHY SELECT GOOD SEEDS




- The larger amount of food in good seeds results in better growth of seedlings.
- Good seeds result in better seedlings – healthier, heavier, and more roots.
- Healthy seedlings will grow faster than the poor seedlings when transplanted in the field.
- Good seeds result in uniform germination and growth.

# SEEDLING GROWTH


- 21 Source of food for growth
- 22 Factors affecting seedling growth — water depth
- 23 Factors affecting seedling growth — amount of water
- 24 Factors affecting seedling growth — temperature
- 25 Factors affecting seedling growth — light intensity
- 26 Factors affecting seedling growth — light intensity
- 27 Factors affecting seedling growth — available nutrients
- 28 Factors affecting seedling growth — available nutrients

# SOURCE OF FOOD FOR GROWTH



- The seedling grows by using food from the endosperm.
- After producing four leaves, it grows from food taken up through the roots and manufactured in the leaves.
- As the seedling gets older, it becomes more dependent on the environment for food.
- A dapog seedling contains very little food in the endosperm at the time of transplanting. It is just becoming independent in manufacturing its own food.


# FACTORS AFFECTING SEEDLING GROWTH — WATER DEPTH



- Deep flooding results in poor root growth and tall seedlings due to lack of air in the soil. When transplanted, the seedlings are easily damaged.




## FACTORS AFFECTING SEEDLING GROWTH— AMOUNT OF WATER




- The large amount of water available to the plants in lowland nursery results in uniform shoot growth.
- The irregular distribution of water in upland nursery results in uneven growth. However, root growth is usually excellent.
- Insufficient water results in slow seedling growth.

# FACTORS AFFECTING SEEDLING GROWTH— TEMPERATURE




- Plants grow faster at warm temperature than at cooler temperatures.
- Seedlings are taller when grown during warm temperatures than during cooler temperatures. Cooler temperatures can cause yellowing of leaves and eventual death of seedlings.

## FACTORS AFFECTING SEEDLING GROWTH— LIGHT INTENSITY



- Seedlings need bright light. Cloudy days mean less brightness.
- Less light means weak seedlings because plants cannot produce enough food.
- Less light can cause elongation of leaf blade and leaf sheath – taller and weaker plants.
- Prepare seedbed away from shadows of trees and buildings.


# FACTORS AFFECTING SEEDLING GROWTH— LIGHT INTENSITY



• Low light intensity results in




— tall and weak seedlings



— seedlings with low dry matter




— seedlings that are easily injured  
at planting




— increased chances of damage  
by diseases

# FACTORS AFFECTING SEEDLING GROWTH— AVAILABLE NUTRIENTS



- Fertilizers supply plant food in addition to what is already available in the soil.
- Fertilizer may be needed if the nursery period is long, in upland nurseries, in cold areas, and in areas with poor soil.

## FACTORS AFFECTING SEEDLING GROWTH— AVAILABLE NUTRIENTS



• Too much fertilizer in the seedbed may result in:

- increased chances of being attacked  
by diseases like blast


- very tall and weak seedling

• Since the nursery period in warm regions of the world is only for 10 to 20 days, use of fertilizer is usually not practiced.

# HOW TO SELECT GOOD SEEDLINGS

- 31 Good seedlings have uniform plant height and growth
- 32 Good seedlings have short leaf sheath
- 33 To have short leaf sheaths use proper water depth
- 34 To have short leaf sheaths good lighting is needed
- 35 Good seedlings have no pests nor diseases such as
- 36 Good seedlings have large number of roots and heavy weight

## GOOD SEEDLINGS HAVE UNIFORM PLANT HEIGHT AND GROWTH




Regular plant growth

Irregular plant growth

- Irregular seedling growth may indicate unevenness in:
  - distribution of seeds on the seedbed
  - germination
  - land preparation of the seedbed
  - watering
  - availability of nutrients in the soil




## GOOD SEEDLINGS HAVE SHORT LEAF SHEATH



- The leaf sheath is the lower portion of the leaf that encloses the stem and young leaves.
- Long leaf sheath indicates very rapid initial elongation, making the seedling weak.


## TO HAVE SHORT LEAF SHEATHS USE PROPER WATER DEPTH



Water level on the seedbed


- Too much water can result in long leaf sheaths and weak seedlings.
- Weak seedlings have poor growth right after transplanting, recovery is slow.
- Long, droopy leaves of poor seedlings tend to stick to the mud when transplanted.

## TO HAVE SHORT LEAF SHEATHS GOOD LIGHTING IS NEEDED




- Low light because of cloudy days, heavy seeding, and shadows from trees can result in long leaf sheaths.


# GOOD SEEDLINGS HAVE NO PESTS NOR DISEASES SUCH AS



Stem borers



Blast



Hoppers


Leaf eaters




Weeds

## GOOD SEEDLINGS HAVE LARGE NUMBER OF ROOTS AND HEAVY WEIGHT

Poor



Good




- Heavy seedlings indicate large amount of food accumulated, resulting in better recovery after transplanting.

# TRANSPLANTING


- 39 Why transplant
- 40 How many seedlings per hill
- 41 Why transplant at the proper depth
- 42 Why cut the leaves of seedlings before transplanting

## WHY TRANSPLANT



- Weed control is simpler in straight row transplanting.
- Direct-seeded rice is susceptible to attack by rats, snails and birds.



# HOW MANY SEEDLINGS PER HILL



- There is no difference in grain yield between one and two seedlings per hill, if no seedlings die.




# WHY TRANSPLANT AT THE PROPER DEPTH



- Tillers normally develop 5 to 10 days after transplanting. Deep planting delays tillering.
- Growth of the plant is set back at transplanting; it takes 2 to 4 days before new roots are formed.

## WHY CUT THE LEAVES OF SEEDLINGS




- Long, droopy leaves of tall seedlings touch the muddy water providing chance of diseases infecting the leaves. Cutting prevents this.
- Wounds caused by cutting may serve as entrance for bacterial infection. To avoid cutting, plant seedlings of right age grown under good conditions.

# THE LEAVES


- 45 The rice leaf
- 46 The leaves of the main stem
- 47 Leaf production
- 48 Internodes

# THE RICE LEAF



- A rice leaf can be distinguished from other grasses by the presence of both the ligule and the auricle.
- A grass leaf has a collar but may have only a ligule or auricle or neither.
- A rice leaf, like the grasses, has parallel veins.


## THE LEAVES OF THE MAIN STEM



- The coleoptile comes out of the seed first, then the primary leaf, the secondary leaf with the first expanded leaf blade, and succeeding leaves.
- The last leaf is called the flag leaf.


# LEAF PRODUCTION

LENGTHWISE SECTION AT THE TIP OF THE STEM



- Rice leaves on the main stem are produced one at a time.
- A new leaf is produced at an average of every 7 days.
- Rice leaves are alternately arranged.

# INTERNODES




- Normally there are four to six elongated (more than 1 cm) internodes at harvest.
- The longer the basal internodes, the greater the tendency for the plant to fall flat on the ground.
- Closer planting, cloudy weather, higher nitrogen level of the soil and higher temperatures will result in longer internodes.

# THE ROOTS

- 51 Origin of roots
- 52 Crown roots
- 53 Root hairs
- 54 Functions of the roots – site of water and nutrient uptake supports the upper parts of the plant
- 55 Root development
- 56 Root development – 30 days after transplanting
- 57 Root development – 50 days after transplanting
- 58 Root development at heading
- 59 Root distribution
- 60 Root distribution depends upon the depth of top soil
- 61 Root distribution depends upon the depth of the plowed layer
- 62 Root distribution depends upon the downward movement of water
- 63 Root distribution depends upon the amount of air available
- 64 Root distribution depends upon placement of fertilizer




## ORIGIN OF ROOTS




- The radicle or primary root usually dies within a month.
- Crown roots develop from the lower nodes.
- Old roots or older parts of a root are colored brown while new and young parts of a root are white.

# CROWN ROOTS




- Crown roots develop from the lower nodes.
- There are two types of crown roots, superficial and ordinary roots.
- Superficial roots develop easily when the air content of the soil is low, as in later growth stage.

# ROOT HAIRS




Cross section of a young root enlarged 120 times




- Root hairs are tubular extensions on the outermost layer of the roots.
- They are important in water as well as nutrient uptake.
- Root hairs are generally short-lived.

# FUNCTIONS OF THE ROOTS – SITE OF WATER AND NUTRIENT UPTAKE SUPPORTS THE UPPER PARTS OF THE PLANT




- Soil water contains nutrients such as nitrogen, phosphorus, and potassium.

# ROOT DEVELOPMENT




- At later stages of plant growth, the roots initiated from the upper nodes develop horizontally and form the superficial or mat roots.

## ROOT DEVELOPMENT— 30 DAYS AFTER TRANSPLANTING




- The roots are found mostly within the plowed layer (18 cm), almost no roots in the subsoil.

# ROOT DEVELOPMENT – 50 DAYS AFTER TRANSPLANTING



- Some roots have grown downward to the subsoil.


## ROOT DEVELOPMENT AT HEADING



- Some big and strong roots have further penetrated into the subsoil.
- Mat roots are abundant.




# ROOT DISTRIBUTION




- Root distribution depends upon:
  - depth of the top soil
  - depth of the plowed layer
  - downward movement of water
  - amount of air available
  - type of irrigation
  - placement of fertilizer
- Roots must penetrate deeply and spread widely and evenly for good uptake of nutrient from the soil.

## ROOT DISTRIBUTION DEPENDS UPON THE DEPTH OF TOP SOIL




- The deeper the top soil or the distance between the surface and the hardpan, the deeper is the root penetration.

# ROOT DISTRIBUTION DEPENDS UPON THE DEPTH OF THE PLOWED LAYER




- The deeper the plowed layer, the deeper the root penetration.
- Plow as deep as possible, shallow plowing restricts root growth.

- If the downward movement of water happens freely and quickly, the roots develop downward easily.
- The downward movement or percolation of water results in more air and fertilizer available in the lower soil layer.
- The deeper the roots the better the water absorbing capacity of the plant. A very important plant characteristic in areas where water supply is not dependable.




## ROOT DISTRIBUTION DEPENDS UPON THE DOWNWARD MOVEMENT OF WATER

## ROOT DISTRIBUTION DEPENDS UPON THE AMOUNT OF AIR AVAILABLE



- Absence of air at a soil layer can result in the decay of roots and inhibition of root development in that layer. A shallow root type develops.
- Downward movement of air dissolved in water depends on the depth and type of top soil.

# ROOT DISTRIBUTION DEPENDS UPON PLACEMENT OF FERTILIZER




- Mixing the fertilizer thoroughly into the plowed soil results in deeper roots and better root distribution.

# THE TILLERS

- 67 Primary tiller
- 68 Tillering pattern
- 69 Production of tillers
- 70 Productive and non-productive tillers
- 71 Percent productive tillers
- 72 Factors affecting tillering – variety
- 73 Factors affecting tillering – spacing
- 74 Factors affecting tillering – season of planting
- 75 Factors affecting tillering – nitrogen level


## PRIMARY TILLER



- The first tiller (primary tiller) usually develops between the main stem and the second leaf from the base.
- Although the tiller is still attached to the mother plant at later stages of growth, it is independent since it produces its own roots.




# TILLERING PATTERN




- Primary (P) tillers come from the main stem.
- Secondary (S) tillers develop from the primary tillers and tertiary (T) tillers from the secondary tillers.
- The lower the point of origin on the main stem, the older is the tiller.

# PRODUCTION OF TILLERS





- Tillering starts 10 days after transplanting and reaches maximum 50 to 60 days after transplanting.
- After reaching the maximum, tiller number decreases as weak tillers die.

## PRODUCTIVE AND NON-PRODUCTIVE TILLERS



- Tillers formed at later stages of growth are usually unproductive. Either the tillers die or the panicles produced are small and too late to catch up with the ripening of the other panicles. The spikelets are only half-filled at the time of harvest.
- Modern varieties have more tillers at flowering and lose less tillers.
- Loss of tillers may be the result of mutual shading, competition among tillers, or lack of nutrients, especially nitrogen!

# PERCENT PRODUCTIVE TILLERS



$$\text{Percent productive tillers} = \frac{\text{number of panicles produced}}{\text{highest number of tillers produced}} \times 100$$

In the above drawings, the percent productive tillers is 50 in the traditional varieties and 75 in the modern varieties.

## FACTORS AFFECTING TILLERING— VARIETY




A variety with 19 tillers




A variety with 54 tillers

- Varieties differ in tillering capacity.
- Tillering capacity can be obtained by growing the plants far apart in rich soil. The full capacity usually is not used under field conditions.

## FACTORS AFFECTING TILLERING — SPACING



50 x 50 cm spacing  
33 tillers per plant  
4 plants per square meter  
122 tillers per square meter




10 x 10 cm spacing  
3 tillers per plant  
100 plants per square meter  
300 tillers per square meter

- The tiller number per plant increases with increase in distance between plants.
- The number of tillers per square meter may be less if plants are spaced too far apart.

## **FACTORS AFFECTING TILLERING— SEASON OF PLANTING**



Rainy season - 21 tillers



Dry season - 16 tillers

- More tillers are produced during the rainy than the dry season.
- More nitrogen fertilizer is needed during dry season to increase tiller number.

## FACTORS AFFECTING TILLERING — NITROGEN LEVEL



10 Tillers  
No nitrogen  
added



30 Tillers  
Fertilizer nitrogen  
added


- The higher the amount of nitrogen added the more tillers produced.



# THE PANICLE

- 79 Panicle formation
- 80 Booting stage
- 81 The spikelet
- 82 Flowering order of a panicle
- 83 Stages of grain formation
- 84 Causes of empty spikelets

# PANICLE FORMATION




- Panicle formation occurs at the tip of the growing point of the shoot.
- The panicle is visible to the naked eye when it is 1 millimeter in length.
- At 1 millimeter, the young panicle has many fine, white, hairy structures at the tip.
- When the panicle inside the leaf sheath is about 1 mm, three more leaves will be produced before the panicle comes out.

## BOOTING STAGE



- At booting stage, there is a bulge at the base of the leaf sheath.
- Booting stage is 20 to 25 days before flowering, the panicle is 1 millimeter in size.
- Flowering occurs 35 days after the start of panicle formation.


# THE SPIKELET



- Opening of the anthers occurs one day after the panicle comes out.
- Low temperature delays the opening of the anthers.
- Pollen from the anthers (like fine dust) have to reach the stigma and unite with the egg inside the ovary for the ovary to develop into a grain.


# FLOWERING ORDER OF A PANICLE

A large panicle with 196 spikelets




- Spikelets on the top branches open first.
- The lower spikelets, which open last, are usually not completely filled in some large panicles.
- Modern varieties have 100 to 120 spikelets per panicle.

## STAGES OF GRAIN FORMATION




- Build-up of starch inside the spikelet begins after the sperm in the pollen unites with the egg in the ovary (fertilization).
- A grain is a ripened ovary together with the lemma and palea.
- At 21 days after fertilization the spikelet has reached its maximum weight.
- Since it takes 7 days for all the spikelets in a panicle to open, full maturity for the whole panicle does not occur until 10 days after flowering.
- Extra days will be needed to ripen all the grains since the panicles do not come out at the same time.

## CAUSES OF EMPTY SPIKELETS



Side view of a fully filled spikelet



Side view of an empty spikelet

- Many factors can affect the filling up of the spikelets such as:
  - lack of starch to fill up the spikelets because of lodging, low light intensity, drying of the leaves, and diseases.
  - drying up of the stigma because of high temperature or dry winds.
  - too much nitrogen applied at panicle formation stage.
  - low temperature and high humidity at flowering resulting in the non-opening of the spikelets.
  - low temperature at panicle formation resulting in degeneration of the pollen.
- Empty spikelets will float when placed in water.


# **DORMANCY**

- 87** Grain dormancy
- 88** Advantages of dormancy – prevents germination of seeds in the panicle
- 89** Advantages of dormancy – prevents germination of seeds if stored in wet conditions for a few days after harvest




# GRAIN DORMANCY

Days after  
harvest



- Dormancy is the failure of a mature seed to germinate under favorable conditions.
- Not all varieties have dormancy.
- Seeds may be dormant from 0 to 80 days depending upon the variety and conditions at harvest.

## ADVANTAGES OF DORMANCY — PREVENTS GERMINATION OF SEEDS IN THE PANICLE



- Dormancy is important during the rainy season harvest.
- When the grains are mature and it rains, the nondormant seeds may germinate.
- Seeds harvested during the dry, sunny season have lower percentage of dormancy.

# ADVANTAGES OF DORMANCY — PREVENTS GERMINATION OF SEEDS IF STORED IN WET CONDITIONS FOR A FEW DAYS AFTER HARVEST



- The cause of dormancy is not clear.
- Dormancy can be a disadvantage, freshly harvested seeds cannot be immediately planted.

# FERTILIZERS


- 93 What is a fertilizer
- 94 Nutrients that the rice plant needs
- 95 Role of fertilizers
- 96 Types of fertilizers – organic
- 97 Types of fertilizers – inorganic
- 98 Fate of nitrogen fertilizer applied to the soil

## WHAT IS A FERTILIZER



- Fertilizers contain important mineral nutrients needed by the plant and are usually applied to the soil.
- Soils sometimes do not provide sufficient nutrients that rice plants need.
- Fertilizers should be applied if mineral nutrients are lacking.

# NUTRIENTS THAT THE RICE PLANT NEEDS



- All of the above except carbon, oxygen, and hydrogen can be supplied by fertilizers.
- There are several mineral nutrients that the plant needs but nitrogen, potassium, and phosphorus are needed in large amounts.

# ROLE OF FERTILIZERS


- manufacture of food



- reproduction



- maintenance of life




- growth



- Nitrogen, phosphorus, and potassium are needed for the life processes going on in the plant.

## TYPES OF FERTILIZERS — ORGANIC

Examples:



- Organic fertilizers come from plant and animal matters such as rotten leaves and chicken manure.
- Large amounts of organic fertilizer contain very small amount of mineral nutrients needed by the plant.
- Use of organic fertilizer results in better soil structure.




# TYPES OF FERTILIZERS— INORGANIC

## Examples:

Urea (45-0-0)


Ammonium sulfate (21-0-0)

Muriate of potash (0-0-60)



- Inorganic fertilizers are commercially manufactured mineral nutrients.
- There are several kinds of combinations of nitrogen, phosphorus, and potassium fertilizers.
- The numbers on the bag refer to the percentage by weight of mineral nutrients in the fertilizer. 24-12-12 means 24% nitrogen, 12% phosphorus ( $P_2O_5$ ), and 12% potassium ( $K_2O$ ).
- The rest of the material in the fertilizer bag is filler material and may contain calcium or sulfur.

## FATE OF NITROGEN FERTILIZER APPLIED TO THE SOIL





- A high percentage of the nitrogen applied is lost.
- Nitrogen fixed in the soil can be partly used by the succeeding crop.
- How to minimize the loss and maximize the use of the available nitrogen is important in good crop management.

# HOW MUCH NITROGEN TO APPLY

- 101 Season of cropping — rainy season
- 102 Season of cropping — dry season
- 103 Fertility of the soil
- 104 Yield potential of the variety
- 105 Profit from fertilizer applied



## SEASON OF CROPPING— RAINY SEASON



Dry season


- Rainy season – plants are tall, leafy, and shade each other so that food manufacturing in the leaves is low. Light energy is low above and inside the crop.
- Fertilizer applied during rainy season cannot be fully used by the plant.
- Lower amount of fertilizer should be used during the rainy season.

## SEASON OF CROPPING — DRY SEASON




- Dry season — plants are shorter and have less tillers, light energy available is more.
- The fertilizer applied increases tiller number, leaf area, and rate of food manufacturing.
- More sunlight and more leaves increase food production — higher profits for fertilizer applied.
- More nitrogen can be applied during the dry season since grain yield as a result of nitrogen application is higher.

## FERTILITY OF THE SOIL




- Too much nitrogen fertilizer in the soil results in too much vegetative growth, resulting in poor distribution of light and possibly lodging of plants.
- Too much nitrogen at later stages of growth increases sterility of spikelets and induces production of late tillers.



- An optimum level of nitrogen in the soil results in the correct leaf area, tiller number and distribution of light, resulting in higher grain yields than the above soil.

# YIELD POTENTIAL OF THE VARIETY


High yield potential – semidwarf



Nitrogen fertilizer

Upright leaves  
increase tillers  
no lodging

Low yield potential – tall




Nitrogen fertilizer

Droopy leaves  
mutual shading  
lodging

- Application of fertilizer to tall varieties will increase their height and tendency to lodge.
- Because of lodging and shading of leaves, grain yields may actually decrease with fertilizer application.

# PROFIT FROM FERTILIZER APPLIED



Low fertilizer  
Low yield  
Low profit

Medium fertilizer  
Medium yield  
High profit

High fertilizer  
Highest yield  
Low profit


- There is a right amount of fertilizer to apply to get maximum profit.
- The right amount of fertilizer will depend on the price of fertilizer in relation to increase in yields.
- The profit from fertilizer applied is higher during the dry than in the rainy season.
- The right amount of fertilizer for high grain yields will vary with the variety.



# HOW TO INCREASE THE EFFICIENCY OF NITROGEN FERTILIZER

- 109 Use high yielding varieties
- 110 Apply the right amount of fertilizer
- 111 Apply fertilizer at correct growth stage of the rice plant
- 112 Prevent the field from drying out
- 113 Mix the fertilizer into the soil
- 114 Do not topdress while the leaves are wet
- 115 Keep the fields free from weeds

# USE HIGH YIELDING VARIETIES




Modern variety




Old variety

## Comparative grain yields




With nitrogen  
modern variety

Dry Season




With nitrogen  
old variety



With nitrogen  
modern variety


Wet Season



With nitrogen  
old variety

- Grain yield increase as a result of nitrogen application is more in the modern than in the old varieties, regardless of the season of planting or amount of nitrogen used.


# APPLY THE RIGHT AMOUNT OF FERTILIZER



● **The right amount of fertilizer will depend upon:**

- season of cropping
- fertility of the soil
- yield potential of the variety
- price of the fertilizer
- time and method of application


# APPLY FERTILIZER AT CORRECT GROWTH STAGE OF THE RICE PLANT



The lighter the shade the better the time of application


- The most efficient times for applying nitrogen fertilizers are at transplanting and at panicle initiation.

## PREVENT THE FIELD FROM DRYING OUT




- On and off flooding results in great losses of nitrogen to the air.
- Nitrogen applied to flooded soil is changed to a different form by the air. This form is easily changed into gas and lost into the air.
- Water keeps the air from moving into the soil. The less air in the soil the less change in nitrogen to gas form. Repair levees to prevent water loss.

## MIX THE FERTILIZER INTO THE SOIL




- Fertilizers applied before transplanting should be mixed thoroughly with the soil:
  - to prevent nitrogen losses to the atmosphere by the action of air.
  - to put the fertilizer nearer to the roots.
- Do not broadcast fertilizer without mixing it into the soil.
- Do not topdress in water immediately after transplanting.

## DO NOT TOPDRESS WHILE THE LEAVES ARE WET



- The fertilizer will stick on the wet leaves and may cause leaf burn.
- The dissolved fertilizer will be lost to the air when the droplets dry up.



- Similarly do not topdress if a heavy rain is impending: The fertilizer might be washed out from the field.

## KEEP THE FIELDS FREE FROM WEEDS





- Weeds compete with the rice plants for the added nitrogen fertilizer.
- Weed before applying nitrogen.
- Weed growth increases with fertilizer application.
- The more vigorous the weed growth the greater will be the competition.



# WHY MORE NITROGEN FERTILIZER IS APPLIED DURING THE DRY SEASON

- 119 Higher grain yields from nitrogen application
- 120 Less danger of shading
- 121 Less danger of lodging
- 122 Increases the low tiller number


# HIGHER GRAIN YIELDS FROM NITROGEN APPLICATION





- There is a greater response to nitrogen fertilizer during the dry season than during the rainy season.
- Sunlight, which is necessary for the manufacture of food, is more abundant during the dry season.

## LESS DANGER OF SHADING


300units



500units




Rainy season



Dry season


- Less leafiness — shorter and more erect leaves during the dry season.
- Less danger of shading since there is sufficient sunlight. The amount of light is higher and leaf arrangement for catching the sunlight is better during the dry season.
- If shading occurs yields are reduced.

## LESS DANGER OF LODGING



- Plants do not grow very tall during the dry season compared with the wet season thus lodging is less likely even with higher rates of nitrogen fertilizer.

## INCREASES THE LOW TILLER NUMBER




- Nitrogen increases the number of tillers.
- Rice generally produces fewer tillers during the dry than in the rainy season.
- One reason for using closer spacing during the dry season is that fewer tillers are formed during this season.
- The additional tillers produced as a result of nitrogen fertilization are mostly productive since there is less shading during the dry season.

# CARBOHYDRATES PRODUCTION


- 125 The food factory
- 126 The food factory
- 127 Factors affecting carbohydrate production –  
amount of green color
- 128 Factors affecting carbohydrate production –  
amount of green color
- 129 Factors affecting carbohydrate production –  
amount of light
- 130 Factors affecting carbohydrate production –  
amount of light
- 131 Factors affecting carbohydrate production –  
amount of water in the leaf
- 132 Factors affecting carbohydrate production –  
amount of air

# THE FOOD FACTORY




- Carbohydrate is manufactured in the green leaves.
- Water from the soil and carbon dioxide from air are the main materials in the manufacture of carbohydrates.
- Water is absorbed by the roots from the soil. Air enters the plants through the leaf pores on the surface of the leaves.

# FACTORS AFFECTING CARBOHYDRATE PRODUCTION— AMOUNT OF GREEN COLOR



Different  
number of leaves




Different  
sizes of leaves


- Total amount of green color (chlorophyll) per plant increases with increase in number and size of the leaves.



## **FACTORS AFFECTING CARBOHYDRATE PRODUCTION— AMOUNT OF GREEN COLOR**




Many manufacturing units



Only few manufacturing units


- Total amount of green color (chlorophyll) per plant increases with increase in thickness of leaves and number of manufacturing units inside the leaves.

# FACTORS AFFECTING CARBOHYDRATE PRODUCTION AMOUNT OF LIGHT



- The brighter the light, the more light energy, resulting in more carbohydrates produced.


## FACTORS AFFECTING CARBOHYDRATE PRODUCTION — AMOUNT OF LIGHT



- There is more light available to plants with erect leaves, thus more carbohydrate is manufactured.

# FACTORS AFFECTING CARBOHYDRATE PRODUCTION—AMOUNT OF WATER IN THE LEAF

Cross section of a highly magnified leaf




- Loss of water content results in closure of pores.
- Water is an important part of a carbohydrate unit.
- Lack of water leads to decreased rate of food manufacture when pores close and air cannot enter.

# FACTORS AFFECTING CARBOHYDRATE PRODUCTION— AMOUNT OF AIR

AMOUNT OF AIR

## FOOD MANUFACTURING =




- The plant uses carbon dioxide from the air for food manufacture.
- Carbon dioxide is plentiful and rarely the cause of a decrease in food manufacturing.
- Thus it can be seen that water, air, light, and green color are needed for food manufacture. If any one of these is lacking then food manufacturing is slowed down even if the others are present in abundance.


# **WATER**

- 135** Major component of the plant
- 136** Raw material for food manufacturing
- 137** Carries the food
- 138** Cools the plant
- 139** Stiffens the plant

# MAJOR COMPONENT OF THE PLANT




# RAW MATERIAL FOR FOOD MANUFACTURING



- Lack of water decreases the amount of food manufacturing.
- Water, air, and light are needed in food manufacturing, water is usually the limiting factor.




# CARRIES THE FOOD



- Water carries the carbohydrates and mineral nutrients to the different plant parts.
- One hectare of rice plants uses at least 8 million liters (400,000 big kerosene cans) of water during its life.

# COOLS THE PLANT

Water - cools the leaf as it evaporates



Cross section of a leaf showing the pores where water evaporates

- Water can cool the leaves similar to perspiration cooling our body.
- Without water in the leaves the pores close: water cannot pass out and air cannot enter. Growth is greatly retarded.
- If the temperature is too high and water does not evaporate, the leaves dry up.
- Most of the water taken up by the rice plant is lost through evaporation.

# STIFFENS THE PLANT

With water

Without water

With water

Without water



Without water the erect  
leaves will droop


Without water the leaves  
will roll

- Water helps in making the leaves erect and fully expanded.
- Water in the plant is like the air in the tires of a car.

# **YIELD COMPONENTS**


- 143** Stages of growth during which yield components are determined
- 144** Stages of growth during which yield components are determined – leaf development and tillering
- 145** Stages of growth during which yield components are determined – panicle formation
- 146** Stages of growth during which yield components are determined – flowering
- 147** Stages of growth during which yield components are determined – ripening
- 148** Variations in yield components
- 149** Importance of yield components
- 150** Importance of yield components
- 151** How to use yield components
- 152** How to use yield components
- 153** How to use yield components
- 154** How to use yield components

# STAGES OF GROWTH DURING WHICH YIELD COMPONENTS ARE DETERMINED




- Every stage of growth contributes to grain yields, good management at all stages is necessary.
- ☞ Environmental factors affect all these stages.

## **STAGES OF GROWTH DURING WHICH YIELD COMPONENTS ARE DETERMINED— LEAF DEVELOPMENT AND TILLERING**



- The number of tillers formed, which determines the number of panicles, is the most important factor in high grain yields.
- Sufficient leaves are necessary to insure large number of spikelets and to fill the spikelets.

## **STAGES OF GROWTH DURING WHICH YIELD COMPONENTS ARE DETERMINED— PANICLE FORMATION**



- The number of spikelets per panicle is determined at this stage.
- Very low temperatures and low available light energy during this stage will increase the number of aborted spikelets.

## **STAGES OF GROWTH DURING WHICH YIELD COMPONENTS ARE DETERMINED — FLOWERING**



- Transfer of the male cell to the egg in the ovary occurs at flowering.
- Successful transfer will determine the development of the spikelet into a grain, the result of accumulation of carbohydrate and formation of the embryo.



## STAGES OF GROWTH DURING WHICH YIELD COMPONENTS ARE DETERMINED - RIPENING



- The weight of a grain is determined at this stage. It is least affected by environmental factors.
- Poor tillering or low tiller number per unit area cannot be compensated by increasing the weight per grain or increasing fertility of the spikelets since both components do not vary much.

## VARIATIONS IN YIELD COMPONENTS



Few but large panicles  
panicle weight type

Many but small panicles  
panicle number type

- Increase in grain yield of panicle *number* types is usually the result of an increase in number of panicles.
- Increase in grain yield of panicle *weight* types is usually the result of an increase in the weight per panicle.
- Most modern high yielding varieties are panicle number types while the traditional varieties are panicle weight types.

# IMPORTANCE OF YIELD COMPONENTS

- Detailed study of the different factors contributing to grain yields can reveal why the yields are high or low.
- Target yield = 80 cavans/hectare or 4,000 kilograms/hectare or 400 grams/square meter  
1 cavan = 50 kilograms of palay
- Characteristics of the variety you are using:
  - number of panicles per hill = 14
  - spikelets per panicle = 100
  - percentage of filled spikelets = 83.3%
  - weight of a single grain = 0.025 grams

# IMPORTANCE OF YIELD COMPONENTS

- To find out the number of panicles you need per hill:

$$\text{Yield} = \begin{array}{c} \text{panicles} \\ \text{per square} \\ \text{meter} \end{array} \times \begin{array}{c} \text{spikelets} \\ \text{per} \\ \text{panicle} \end{array} \times \begin{array}{c} \text{percent} \\ \text{filled} \\ \text{spikelets} \end{array} \times \begin{array}{c} \text{weight} \\ \text{of a single} \\ \text{grain} \end{array}$$

$$400 \text{ g} = (\text{panicles/sq m}) \times (100) \times \frac{83.3}{100} \times (0.025)$$

$$\begin{aligned} \text{Panicles/sq m} &= \frac{400}{100 \times 0.833 \times 0.025} \\ &= 192 \end{aligned}$$

- If the spacing you used was 25 × 25 cm or 16 hills/sq m

$$\frac{192 \text{ panicles/sq m}}{16 \text{ hills/sq m}} = 12 \text{ panicles/hill}$$

- The variety you are using can produce more than 12 panicles per hill at 25 × 25 cm spacing. So, target yield could be obtained.


- If yield did not meet target:

If the yield actually obtained was below 400 grams per square meter although you were using the correct variety and spacing, something was wrong with your crop. A detailed study of the yield components may reveal what was possibly wrong during the growth of the plants.

# HOW TO USE YIELD COMPONENTS


THE PROBLEM :

Expected



14 panicles

Actual




6 panicles

• What was wrong: probably a defect in the soil or in application of fertilizer. It also could have been lack of water or damage by pests and diseases during earlier growth.

# HOW TO USE YIELD COMPONENTS


The problem:



- What was wrong: the problem occurred a little before, during, or after the formation of the spikelet (26 to 16 days before flowering). It possibly resulted from lack of sunlight, lack of food, or insect damage to the leaves.

# HOW TO USE YIELD COMPONENTS


The problem:



- What was wrong: maybe the temperature was too low ( $20^{\circ}\text{C}$ ) or too high (above  $35^{\circ}\text{C}$ ), the plants lodged, or suffered from lack of water at flowering time. Maybe too much nitrogen was applied.


# HOW TO USE YIELD COMPONENTS

Expected



25 grams per 1000 grains

Actual



20 grams per 1000 grains

- What was wrong: conditions after flowering were unfavorable such as not enough food, not enough leaves to manufacture the food, or cloudy weather.



# **PLANT TYPE OF A LOWLAND RICE VARIETY WITH HIGH GRAIN YIELD POTENTIAL**



- 157 Short stature
- 158 Non-lodging
- 159 Good distribution of light
- 160 Erect leaves
- 161 Flag leaf higher than the panicle
- 162 Short leaves
- 163 Good tillering
- 164 Erect tillers
- 165 Ideal tiller

## SHORT STATURE




- Reduction in plant height is the most important factor in increasing the grain yield potential of rice.
- Reduction in plant height increases resistance to lodging.

# NON-LODGING





- Plant height increases with nitrogen application and lodging becomes a problem.
- Many leaves on the lodged plants decay since they are soaked in water and do not receive sufficient light.
- Short, stiff stem prevents lodging.

# GOOD DISTRIBUTION OF LIGHT



- Tall, leafy plants result in very little light received by the lower leaves.
- Upright tillers and leaves reaching above the panicles contribute to better light distribution, resulting in better food manufacturing and grain yields.


# ERECT LEAVES



Very little light is received by the lower leaves


Very little shading of lower leaves, some spacing and leaf length

# FLAG LEAF HIGHER THAN THE PANICLE




- There is less shading of the upper leaves if the panicle does not extend far above the flag leaf.

## SHORT LEAVES



- Shorter leaves are more erect because they have less weight to carry.
- Erect leaves allow more light to reach the lower parts of the plants.

## GOOD TILLERING




Poor  
tillering  
ability

Good  
tillering  
ability


- Good tillering ability insures adequate tillers per unit area even if some plants die at early stage of growth.



# ERECT TILLERS




Spreading tillers



Erect tillers

- Upright tillers result in better light distribution.

# IDEAL TILLER



● Main tiller at flowering.

# **FACTORS AFFECTING LODGING**

- 169** Plant height
- 170** Method of sowing
- 171** Type of leaf sheath
- 172** Stem thickness
- 173** Wind and rain
- 174** Light intensity
- 175** Spacing
- 176** Amount of fertilizer

## PLANT HEIGHT




Not resistant



Resistant


- The taller the plant the greater is the tendency to lodge.
- Avoid using tall varieties during the rainy season.

## METHOD OF SOWING




- Transplanted rice is more resistant to lodging because the base of the plant is better anchored.

# TYPE OF LEAF SHEATH




# STEM THICKNESS



- The thicker the stem and the thicker the internode, the higher the resistance to lodging.


# WIND AND RAIN



- Wind and rain can lodge a plant. The stronger the wind the more likely the plant will lodge.
- Avoid using tall varieties during the rainy season.



# LIGHT INTENSITY



Dry season



Plants shorter


Rainy season




Plants taller

- Cloudy weather results in taller plants, hence greater tendency to lodge.
- Lodging is more common during the rainy season.

# SPACING





Spacing is too close



Correct spacing

- Close spacing results in taller plants and weaker stems.

# AMOUNT OF FERTILIZER




- Fertilizer increases plant height. Tall varieties cannot stand too much fertilizer.

# WEEDS


- 179 Weeds reduce rice yields
- 180 Weeds compete with rice
- 181 Weeds decrease the effect of nitrogen fertilizer
- 182 Weeds – differences between grasses, sedges, and broadleaved weeds
- 183 Common weeds in rice fields – grass
- 184 Common weeds in rice fields – sedge
- 185 Common weeds in rice fields – broadleaf
- 186 Differences between grasses and rice plants
- 187 When to weed the rice crop

# WEEDS REDUCE RICE YIELDS

Grain yield during the dry season




Weeded




Not weeded

Grain yield during the wet season




Weeded



Not weeded

● Weeds reduce grain yields regardless of the season of planting.


## WEEDS COMPETE WITH RICE



- Weeds compete with rice for sunlight, nutrients, and water.
- If any of these is lacking the others cannot be used effectively even if present in large amounts.
- The competition results in poor rice growth, thus less grain yield.








# WEEDS DECREASE THE EFFECT OF NITROGEN FERTILIZER

Grain yield



- The higher the amount of nitrogen applied, the less is the grain yield if the crop is not weeded.
- Weeds compete with rice for the applied nitrogen fertilizer.
- Nitrogen application favors the growth of weeds more than the rice crop.
- Nitrogen fertilizers should not be used before weeds are controlled.

# WEEDS – DIFFERENCES BETWEEN GRASSES, SEDGES AND BROADLEAVED WEEDS

Type	Grasses	Sedges	Broadleaves
Leaf shape			
Vein arrangement			
Stem cross section			
Examples	Echinochloa crus-galli (Bayakibok)	Cyperus rotundus (Mutha)	Monochoria vaginalis (Gabing uwak)



## COMMON WEEDS IN RICE FIELDS— GRASS



Scientific name: *Echinochloa crusgalli*

Common name: Bayakibok

## COMMON WEEDS IN RICE FIELDS — SEDE



Scientific name: *Cyperus iria*


Common name: Sud-sud

## COMMON WEEDS IN RICE FIELDS — BROADLEAF




Scientific name: *Monochoria vaginalis*  
Common name: Gabing uwak


# DIFFERENCES BETWEEN GRASSES AND RICE PLANTS



Rice – with ligule and auricle




Grass – no ligule, no auricle



Grass – with ligule, no auricle

# WHEN TO WEED THE RICE CROP

Grain yield



Weed-free days after transplanting

- Weeding in the first 30 days following transplanting of the rice plant is important.
- Grain yield is drastically reduced if rice is not weeded during the early stages of growth.

# **CONTROL OF WEEDS**

- 191** Weeds can be controlled by hand pulling
- 192** Weeds can be controlled by mechanical means
- 193** Weeds can be controlled by proper water management
- 194** Weeds can be controlled by proper land preparation
- 195** Weeds can be controlled by crop competition
- 196** Weeds can be controlled by herbicides

# WEEDS CAN BE CONTROLLED BY HAND PULLING

BROADLEAF



Scientific name *Monochoria vaginalis*  
Common name Gabing Uwak

- Pulling weeds by hand is a manual method of control.
- Hand pulling is time consuming.


## **WEEDS CAN BE CONTROLLED BY MECHANICAL MEANS**



- Use of rotary weeder is more efficient than hand weeding.
- Straight row planting is necessary if rotary weeder is used.
- Standing water should be drained from the field when the rotary weeder is used.



## WEEDS CAN BE CONTROLLED BY PROPER WATER MANAGEMENT



Growth of weeds greatly reduced

Growth of weeds slightly reduced

- Most grasses and sedges will be prevented from growing when covered with 5–10 cm water.
- Some broadleaved weeds are not controlled by flooding.
- Many weed seeds do not germinate under water.

## WEEDS CAN BE CONTROLLED BY PROPER LAND PREPARATION



Unevenly prepared land

- Weed growth is encouraged when land is poorly and unevenly prepared and areas are not covered by water.

## WEEDS CAN BE CONTROLLED BY CROP COMPETITION



10 x 10 cm spacing



15 x 15 cm spacing




20 x 20 cm spacing

- The closer the plant spacing the fewer the weeds — less light for the weeds to germinate and grow.
- The shorter the weeds the lesser is the weed damage.

# WEEDS CAN BE CONTROLLED BY HERBICIDES

Applying powders or liquids in solution




Applying granules

# HERBICIDES


- 199 Types of herbicides based on formulation
- 200 Types of herbicides based on time of application
- 201 Types of herbicides based on selectivity
- 202 Types of herbicides based on types of action
- 203 Rice injuries from too much herbicide – tillers tend to spread out
- 204 Rice injuries from too much herbicide – occurrence of brown spots
- 205 Rice injuries from too much herbicide – formation of onion-like leaves
- 206 Rice injuries from too much herbicide – dwarfing of the plant
- 207 Herbicides may kill plants by preventing the manufacture of food
- 208 Herbicides may kill plants by interfering with the plant system

# TYPES OF HERBICIDES BASED ON FORMULATION



- Commercial herbicides are available in powder, liquid or granular forms.
- Granular forms are broadcast and no special equipment needed for application.


## TYPES OF HERBICIDES BASED ON TIME OF APPLICATION



Before the weed seedlings  
come out.

After the weed seedlings  
are out.

# TYPES OF HERBICIDES BASED ON SELECTIVITY




Selective herbicides will kill certain plants only at low concentrations.


Non-selective herbicides will kill all plants.



# TYPES OF HERBICIDES BASED ON TYPES OF ACTION



SYSTEMIC ( TRANSLOCATED )



- Contact herbicides will kill only the plant parts that were sprayed.
- Systemic (translocated) herbicides can travel inside the plant and can therefore kill the whole plant.


# RICE INJURIES FROM TOO MUCH HERBICIDE— TILLERS TEND TO SPREAD OUT



Sprayed too much  
herbicide

Sprayed the correct  
amount of herbicide

## RICE INJURIES FROM TOO MUCH HERBICIDE— OCCURRENCE OF BROWN SPOTS




Too much

Right amount

- Herbicide injury may look like a blast or cercospora leaf spot but closer look shows that the spots are discretely round.

# RICE INJURIES FROM TOO MUCH HERBICIDE— FORMATION OF ONION-LIKE LEAVES



- The new leaves coming out are tube-like or cylindrical if the amount of herbicide applied was too great.

## RICE INJURIES FROM TOO MUCH HERBICIDE— DWARFING OF THE PLANT




Too much

Right amount


- Be sure to use the correct amount of herbicide. Follow the recommended rate.

# HERBICIDES MAY KILL PLANTS BY PREVENTING THE MANUFACTURE OF FOOD



- Herbicides can stop the many different activities going on in the rice plant.
- The manufacture of food involves many steps. A herbicide can block one or more of these steps.

## HERBICIDES MAY KILL PLANTS BY INTERFERING WITH THE PLANT SYSTEM




- The manufacture of protein and production of energy leading to plant growth involves hundreds of steps.
- A different protein-compound is responsible for each step. An herbicide might damage any of these protein compounds.
- Any break in the steps may cause death to the plant.

# HOW TO JUDGE A RICE CROP AT FLOWERING

- 211 At flowering a good rice crop should have uniform plant height
- 212 At flowering a good rice crop should have no lodging
- 213 Lodging may indicate spacing used was too close
- 214 Lodging may indicate fertilizer applied was too much
- 215 Lodging may indicate variety used was too tall
- 216 At flowering a good rice crop should have white to brown roots
- 217 At flowering a good rice crop should have green and undamaged leaves
- 218 At flowering a good rice crop should have at least 3 to 4 leaves per tiller
- 219 At flowering a good rice crop should have the correct plant density
- 220 At flowering a good rice crop should have 250 to 350 panicles per square meter




## **AT FLOWERING A GOOD RICE CROP SHOULD HAVE UNIFORM PLANT HEIGHT**



- Irregular plant height can mean many things:
  - plants lacked water or suffered from stem borer attack or virus diseases.
  - uneven land preparation.
  - uneven fertilization.
  - the seeds used were not pure.

# AT FLOWERING A GOOD RICE CROP SHOULD HAVE NO LODGING



Lodged




Upright

- Lodging may indicate:
  - spacing used was too close
  - too much fertilizer was applied
  - variety used was too tall for that area and for the season of planting.

# LODGING MAY INDICATE SPACING USED WAS TOO CLOSE




Spacing too close




Correct spacing

## LODGING MAY INDICATE FERTILIZER APPLIED WAS TOO MUCH



Too much




Proper amount

- Too much fertilizer causes plant to grow too tall, then it lodges.

# LODGING MAY INDICATE VARIETY USED WAS TOO TALL



Lodged




Upright

## AT FLOWERING A GOOD RICE CROP SHOULD HAVE WHITE TO BROWN ROOTS



- Black roots and foul smell indicate something is wrong with the soil:
  - lack of drainage
  - iron toxicity
  - lack of air
  - high organic acids

## AT FLOWERING A GOOD RICE CROP SHOULD HAVE GREEN AND UNDAMAGED LEAVES




Undamaged leaf

Damaged leaves

- Green undamaged leaves indicate no deficiency or toxicity in the soil, and no attacks by pests or diseases.
- Yellow leaves may indicate nitrogen deficiency or virus diseases.


## AT FLOWERING A GOOD RICE CROP SHOULD HAVE AT LEAST 3 TO 4 LEAVES PER TILLER



- A tiller needs 3 to 4 leaves:
  - to better provide the roots and other plant parts with sufficient food.
  - to better fill up the spikelets with starch manufactured in the leaves.
- If a tiller has only 2 leaves, suspect some deficiency in the soil or water stress at an earlier stage of growth.




## AT FLOWERING A GOOD RICE CROP SHOULD HAVE THE CORRECT PLANT DENSITY




- The correct number of plants per unit area can be checked by standing on the levee. If one can barely see the water or sparkle of the sun rays, then the plant density is right.
- If one cannot see the water, probably the spacing was too close, too much fertilizer was applied, or the variety was too tall.

# AT FLOWERING A GOOD RICE CROP SHOULD HAVE 250 TO 350 PANICLES PER SQUARE METER

Start counting here



- Count the number of panicles per hill (clump) in at least 3 hills inside the field. Do not count the first 3 rows from the levee.
- Get the spacing used.



If the distance between hills is: 25 X 25 cm

Area per hill = 25 X 25 = 625 sq m = 0.0625 sq m

$$\begin{aligned} \text{Number of hills per sq m} &= \frac{1 \text{ sq m}}{\text{area per hill}} \\ &= \frac{1}{0.0625} \\ &= 16 \end{aligned}$$

- To get the number of panicles per square meter

Assuming: 17 panicles per hill  
16 hills per sq m

$$\begin{aligned} \text{Number of panicles per sq m} &= \text{Number of panicles per hill} \times \text{Number of hills per sq m} \\ &= 17 \times 16 \\ &= 272 \end{aligned}$$

- If number of panicles per square meter is less than 250 something is wrong with the method of farming, the rice variety or the soil. Check spacing and fertilizer application.